

公開日： 2020年4月

プロトコール名	1クール	術前	3週	切除不能・再発	4週
フルオロウラシル・ランダ療法 (プロイメンド) (術前・切除不能・再発食道癌) 外科	総クール	2クール			未定
	休薬期間	2週			3週

番号	薬剤名・用量(mg/body)	術前																											
		切除不能進行再発																											
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
①	生食 500mL	→																											
②	生食 500mL + 硫酸Mg補正液 0.4A	→																											
③	生食50mL		→	→	→	→																							
④	フルオロウラシル(800mg/m ²) + ソルデム1 500mL	→	→	→	→	→																							
⑤	生食50mL		→	→	→	→	→																						
⑥	プロイメンド(150) 1V + 生食 100mL (④と同時に開始)	→																											
⑦	生食 100mL + アロキシ 1V + テキサート(1.65) 6A	→																											
⑧	ランダ(80mg/m ²) + 生食(total500mL)	→																											
⑨	フロセミド(20) 1A	→																											
⑩	生食 100mL + テキサート(1.65) 4A		→	→	→	→																							
⑪	生食500mL	→	→	→	→	→																							
⑫	生食500mL	→	→	→	→	→																							
⑬	生食500mL		→	→	→	→																							