

公開日： 2026年8月

プロトコール名 イミフィンジ・GC療法 (Ccr≥60) (尿路上皮癌) 泌尿器科	1クール	術前化学療法	2投1休
		術後化学療法	1投3休
	総クール	術前化学療法	最大4サイクル
		術後化学療法	最大8サイクル
	休薬期間	術前化学療法	1週間
		術後化学療法	3週間

術後化学療法																						
番号	薬剤名・用量 (mg/body)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
①	生食50mL	→																				
②	イミフィンジ(1500mg/body) + 生食250mL	→																				
③	生食50mL	→																				
④	グラニセロン3mgバッグ 1袋 + デキサート(1.65) 4A	→							→													
④'	ハロノセロン0.75mgバッグ 1袋 + デキサート(1.65) 4A		→																			
⑤	ケムシタビン(1000mg/m ²) + 5%糖液 100mL	→							→													
⑥	生食50mL	→							→													
⑦	ソルデム3A 500mL		→	→																		
⑧	ソルデム1 500mL		→																			
⑨	マンニトール 300mL		→																			
⑩	ランダ(70mg/m ²) + 生食 (total 500mL)		→																			
⑪	マンニトール 300mL		→																			
⑫	ソルデム3A 500mL		→																			
⑬	ソルデム1 500mL		→																			

術後化学療法																													
番号	薬剤名・用量(mg/body)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
①	生食50mL	→																											
②	イミフィンジ(1500mg/body) + 生食250mL	→																											
③	生食50mL	→																											