

公開日： 2024年5月

プロトコール名	1クール	3週間
Pola-BR療法 (再発・難治 びまん性大細胞型 B細胞リンパ腫) 内科	総クール	未定
	休薬期間	2週間

1クール目		0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
番号	薬剤名・用量 (mg/body)																						
①	アセリオ静注1000mg バッグ 0.5P	→	→																				
②	ホラミン(5)1A+ソル・マトロール(125)1V+生食100mL	→	→																				
③	生食50mL	→																					
④	リツキシマブ(375mg/m ²) + 生食(10倍希釈)	→																					
⑤	生食50mL	→																					
⑥	ポライビー(1.8mg/kg) + 生食100mL		→																				
⑦	パロノセトロン0.75mg/バッグ1P		→																				
⑦'	パロノセトロン0.75mg/バッグ1P + デキサート(1.65) 4A			→																			
⑧	トレアキシシン(90mg/m ²)+ 生食50mL		→	→																			
⑨	生食50mL		→	→																			
⑩	ジーラスタ皮下注 3.6mg				→																		

2クール以降		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	
番号	薬剤名・用量 (mg/body)																						
①	アセリオ静注1000mg バッグ 0.5P	→																					
②	ホラミン(5)1A+ソル・マトロール(125)1V+生食100mL	→																					
③	生食50mL	→																					
④	リツキシマブ(375mg/m ²) + 生食(10倍希釈)	→																					
⑤	生食50mL	→																					
⑥	ポライビー(1.8mg/kg) + 生食100mL	→																					
⑦	パロノセトロン0.75mg/バッグ1P	→																					
⑦'	パロノセトロン0.75mg/バッグ1P + デキサート(1.65) 4A		→																				
⑧	トレアキシシン(90mg/m ²)+ 生食50mL	→	→																				
⑨	生食50mL	→	→																				
⑩	ジーラスタ皮下注 3.6mg			→																			