

公開日: 2024年7月

プロトコール名 イミフィンジ・GC療法 (胆道癌) 消化器内科	1クール	3週間(9クール目 以降4週間)
	総クール	未定
	休薬期間	1週間(9クール目 以降3週間)

1~8クール目		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
番号	薬剤名・用量(mg/body)																					
①	生食50mL	→																				
②	イミフィンジ + 生食 250mL	→																				
③	生食50mL	→																				
④	NS 500mL	→							→													
⑤	NS 500mL	→							→													
⑥	NS 500mL	→							→													
⑦	ホスアプレピタント(150) + 生食100mL	→							→													
⑧	パロノセトロンバッグ0.75mg 1P + デキサート(1.65)4A	→							→													
⑨	ランダ(25mg/m ²) + 生食(total250mL)	→							→													
⑩	ゲムシタビン(1000mg/m ²) + 5%糖液100mL	→							→													
⑪	生食50mL	→							→													

9クール目以降		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26
番号	薬剤名・用量(mg/body)																										
①	生食50mL	→																									
②	イミフィンジ + 生食 250mL	→																									
③	生食50mL	→																									

番号	薬剤名・用量 (mg/body)	27	28
①	生食50mL		
②	イミフィンジ + 生食 250mL		
③	生食50mL		