

公開日: 2024年6月

プロトコール名 イミフィンジ・GC療法 (胆道癌) 腫瘍内科	1クール	3週間(9クール目 以降4週間)
	総クール	未定
	休薬期間	1週間(9クール目 以降3週間)

1~8クール目

番号	薬剤名・用量(mg/body)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	
①	生食50mL	→																					
②	イミフィンジ + 生食 250mL	→																					
③	生食50mL	→																					
④	NS 500mL	→							→														
⑤	NS 500mL	→							→														
⑥	NS 500mL	→							→														
⑦	ホスアプレピタント(150) + 生食100mL	→							→														
⑧	パロノセトロンバッグ0.75mg 1P + デキサート(1.65)4A	→							→														
⑨	ランダ(25mg/m <sup>2</sup> ) + 生食(total250mL)	→							→														
⑩	ゲムシタビン(1000mg/m <sup>2</sup> ) + 5%糖液100mL	→							→														
⑪	生食50mL	→							→														

9クール目以降

番号	薬剤名・用量(mg/body)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26
①	生食50mL	→																									
②	イミフィンジ + 生食 250mL	→																									
③	生食50mL	→																									

番号	薬剤名・用量 (mg/body)	27	28
①	生食50mL		
②	イミフィンジ + 生食 250mL		
③	生食50mL		